

### Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 28 January 2013

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Tuesday, January 22, 2013



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s). Cell concentration sampling data from January 20 to 24 shown as red (high), orange (medium), yellow (low b), brown (low a), blue(very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs\_bulletin\_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at: http://www.tpwd.state.tx.us./landwater/water/environconcerns/hab/redtide/status.phtml

http://tidesandcurrents.noaa.gov/hab/bulletins.html


# **Conditions Report**

There is currently no indication of a harmful algal bloom of Karenia brevis (commonly known as Texas red tide) at the coast in Texas. No respiratory impacts are expected alongshore Texas today through Monday, February 4. For information on area shellfish restrictions, contact the Texas Department of State Health Services.

## Analysis

There is currently no indication of a harmful algal bloom of *Karenia brevis* at the coast in Texas. Recent MODIS imagery (1/26) indicates elevated to very high (3 to > 10  $\mu$ g/L) chlorophyll levels stretching along- and offshore from Sabine Pass almost to the Matagorda Peninsula, with elevated chlorophyll (2-7  $\mu$ g/L) visible along- and offshore from Matagorda Island to the Rio Grande. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast. Forecast models based on predicted near-surface currents indicate a potential transport of 10 km north from the Port Aransas region from January 26-31.

#### Urizar, Kavanaugh




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

#### Wind Analysis

**Port Aransas**: Southeasterly winds (15-20 kn, 8-10 m/s) today. Southerly to southwesterly winds (15-25 kn, 8-13 m/s) Tuesday shifting to northwesterly winds after midnight. Northwesterly winds (15-25 kn) Wednesday. Northerly to northeasterly winds (10-20 kn, 5-10 m/s) Wednesday night. Southeasterly winds (5-15 kn, 3-8 m/s) Thursday. Easterly winds (5-15 kn) Friday and southeasterly winds (10-15 kn, 5-8 m/s) Friday night.

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive:



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).